



Chelation Blend™



Clinical Applications

- Supports Removal of Body Stores of Heavy Metals
- Binds to Metallic Elements and Assists in the Maintenance of a Normal Balance of Metal Elements in the Cells
- Supports Ongoing Chelation Between Intravenous Chelation Treatments

Chelation Blend™ is an extraordinary all-natural combination of unique and therapeutic ingredients formulated to support the safe and effective removal of heavy metals. It is designed to protect against heavy metal-induced free radical damage with the addition of the powerful antioxidants, N-Acetyl-Cysteine and Alpha Lipoic Acid.

All Chantilly Natural Health® Formulas Meet or Exceed cGMP quality Standards

Discussion

DMSA is a non-toxic, water soluble 2,3-dimercaptosuccinic acid employed as a metal chelator since the 1950's. Containing two sulfhydryl groups, this relatively stable compound with high affinity for the toxic metal, penetrates into tissue deposits of the metal, only minimally reacts with essential biological molecules and rapidly eliminates the chelated toxic metal through urine and stool.¹

EDTA, Ethylene diamine tetra-acetic acid is a weak, synthetic amino acid and perhaps the most well known and often used chelating agent. EDTA has the ability to bind with virtually every metal in the Periodic Table.² EDTA is FDA-approved for lead, mercury, aluminum and cadmium toxicity. There is benefit in combining EDTA with other chelators because alone it acts slowly and could potentially lead to redeposition of metals. It is important to replace beneficial minerals such as zinc, copper, iron, cobalt and manganese that are bound by the EDTA.³

Cilantro (*Coriandrum sativum*) (Chinese parsley, coriander) suppresses the deposition of lead by chelating the metal.⁴ A sorbent prepared from cilantro was found to have good efficiency in removing organic and methyl mercury from aqueous solutions.⁵

Chlorella, green algae has demonstrated the ability to take up toxic metals in its environment decreasing resorption of heavy metals excreted into the gut from the liver-biliary system. Chlorella's influence is synergistic with that of garlic and cilantro. Researchers observed a dramatic reduction of 66.03% in blood lead levels in a group of mice receiving chlorella extract concurrent with lead exposure.⁶

Garlic in a variety of forms has been shown by researchers to be beneficial in the management of heavy metal toxicity especially lead,⁷ cadmium⁸ and mercury.⁹

N-Acetyl-L-Cysteine (NAC) has the ability to interact with reactive oxygen species (ROS) and stimulate the body to produce glutathione enhancing cell survival after exposure to heavy metals or toxins.¹⁰

Alpha Lipoic Acid is mostly known for its attribute as a powerful antioxidant. However, it may also directly chelate heavy metals especially copper,¹¹ arsenic,¹² cadmium¹³ and to a small degree, mercury.¹⁴



Supplement Facts

Serving Size: 4 Capsules
Servings Per Container: 30

	Amount	% Daily Value
DMSA	100 mg	**
EDTA (as calcium EDTA)	300 mg	**
Cilantro	300 mg	**
Chlorella	300 mg	**
Allicin (as organic garlic extract)	6,000 mcg	**
N-Acetyl Cysteine	200 mg	**
Alpha Lipoic Acid	100 mg	**

** Daily value not established

Other Ingredients: HPMC, Magnesium Stearate, and Stearic Acid

Dosing:

Professional guidance is recommended, as serving size varies according to individual need (health status). May be used to enhance IV protocols, though not formulated to replace IV chelation. Prophylactic use, e.g., after seafood consumption, take 1 or 2 capsules directly after meal.

References

1. Aaseth J. Recent advance in the therapy of metal poisonings with chelating agents. *Hum Toxicol.* 1983 Apr;2 (2):257-72. [PMID: 6345341] [Harvey, F. Xymogen Xtravaganza presentation, Orlando, FL; June, 2004]
2. Bothwell TH, MacPhail AP. The potential role of NaFeEDTA as an iron fortificant. *Int J Vitam Nutr Res.* 2004 Nov;74(6):421-34 [PMID: 15743018]
3. Gurer H, Ercal N. Can Antioxidants Be Beneficial in the Treatment of Lead Poisoning?, *Free Radic Biol Med*, 2000;29(10): 927-945.[Clinical Pearls newsletter, Sacramento, CA, 2001]
4. Aga M, et al. Preventive effect of *Coriandrum sativum* (Chinese parsley) on localized lead deposition in ICR mice. *J Ethnopharmacol.* 2001 Oct;77(2-3):203-8. [PMID: 11535365]
5. Karunasagar D, Krishna MV, Rao SV, Arunachalam J. Removal and preconcentration of inorganic and methyl mercury from aqueous media using a sorbent prepared from the plant *Coriandrum sativum*. *J Hazard Mater.* 2005 Feb 14;118(1-3):133-9. [PMID: 15721537]
6. Queiroz ML, et al. Protective effects of *Chlorella vulgaris* in lead-exposed mice infected with *Listeria monocytogenes*. *Int Immunopharmacol.* 2003 Jun;3(6):889-900. [PMID: 12781705]
7. Senapati SK, Dey S, Dwivedi SK, Swarup D. Effect of garlic (*Allium sativum* L.) extract on tissue lead level in rats. *J Ethnopharmacol.* 2001 Aug;76(3):229-32 [PMID: 11448543]
8. Jiang W, Liu D, Hou W. Hyperaccumulation of cadmium by roots, bulbs and shoots of garlic (*Allium sativum* L.). *Bioresour Technol.* 2001 Jan;76(1):9-13 [PMID: 11315815]
9. Lee JH, Kang HS, Roh J Protective effects of garlic juice against embryotoxicity of methylmercuric chloride administered to pregnant Fischer 344 rats. *Yonsei Med J.* 1999 Oct;40(5):483-9. [PMID: 10565261]
10. James SJ, Slikker W 3rd, Melnyk S, New E, Pogribna M, Jernigan S. Thimerosal neurotoxicity is associated with glutathione depletion: protection with glutathione precursors. *Neurotoxicology.* 2005 Jan;26(1):1-8 [PMID: 15527868]
11. Gaetke LM, Chow CK. Copper toxicity, oxidative stress, and antioxidant nutrients. *Toxicology.* 2003 Jul 15;189(1-2):147-63. [PMID: 12821289]
12. das Neves RN, Carvalho F, Carvalho M, Fernandes E, Soares E, de Bastos ML, de Pereira ML. Protective activity of hesperidin and lipoic acid against sodium arsenite acute toxicity in mice. *Toxicol Pathol.* 2004 Sep-Oct;32 (5): 527-35 [PMID: 15603538]
13. Bludovska M, Kotyzova D, Koutensky J, Eybl V. The influence of alpha-lipoic acid on the toxicity of cadmium. *Gen Physiol Biophys.* 1999 Oct;18 Spec No:28-32. [PMID: 10703716]
14. Keith RL, Setiarahardjo I, Fernando Q, Aposhian HV, Gandolfi AJ. Utilization of renal slices to evaluate the efficacy of chelating agents for removing mercury from the kidney. *Toxicology.* 1997 Jan 15;116(1-3):67-75 [PMID: 9020508]

Precautions:

Supervise closely. Insufficient data to state safety in pregnancy, lactation, children, elderly or for individuals on drug therapy. Be sure patient has good mineral status before using Chelation Blend™; replete minerals (MinRx™) after the Chelation Blend™.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.